

Abstract 323811: Impact of treatment sequence on survival outcome in patients with a 2nd treatmentline for mCRPC: A new-user design in the French nationwide claims database

M. Gross-Goupil¹, N. Thurin^{2*}, M. Rouyer², T. Haaser¹, X. Rébillard³, M. Soulié⁴, G. de Pouvourville⁵, F. Messaoudi⁶, C. Capone⁶, E. Bignon², J. Jové², S. Lamarque², C. Droz-Perroteau², N. Moore^{1, 2}, P. Blin²

¹ CHU de Bordeaux, Bordeaux, France; ² Bordeaux PharmacoEpi, INSERM CIC1401, Université de Bordeaux, Bordeaux, France; ³ Clinique Beau Soleil, Montpellier, France; ⁴ CHU de Toulouse, Toulouse, France; ⁵ ESSEC, Cergy-Pontoise, France; ⁶ Marketing authorisation holder of abiraterone acetate

BACKGROUND

Abiraterone acetate in association with prednisone/prednisolone, and docetaxel can both be used as 1st or 2nd line treatments for metastatic castration-resistant prostate cancer (mCRPC).

Is it better to start with an abiraterone acetate 1st-line followed by a docetaxel 2nd-line (ABI-DOCE sequence) or to use the inverse sequence DOCE-ABI?

METHODS

- mCRPC patients were identified in the **French National Healthcare System database (SNDS)** using a validated algorithm
- SNDS covers the French population from birth to death and includes out and inpatients information
- To be included, patients had:
 - ✓ To be **aged ≥40** and **covered by the Régime Général** health insurance (86% of the French population)
 - ✓ To **have initiated in 2014** an abiraterone acetate 1st-line followed by a docetaxel 2nd-line (**ABI-DOCE sequence**) or a docetaxel 1st-line followed by an abiraterone acetate 2nd-line (**DOCE-ABI sequence**), all drugs presumed to be used according to the Summary of Product Characteristics
 - ✓ To have a **3-year follow-up** and **5-year history** with no gap >1 year

- A **high dimensional propensity score** (hdPS), was calculated for each patient of each cohort: estimation of the probability for a patient to be treated by ABI-DOCE sequence *versus* DOCE-ABI sequence based on forced and empirically selected variables from 5 dimensions:

Forced variables	Dimensions for variable empirical selection
<ul style="list-style-type: none">Age at index dateCancer stage prior to mCRPC statusCharlson comorbidity index	<ul style="list-style-type: none">Long term disease registrationHospital discharge diagnosesDispensed drugsPerformed laboratory testsPerformed medical procedures

- Patients were 1:1 matched** on hdPS (+/- 0.01), cancer stage prior to mCRPC and date of initial diagnosis (+/- 1 year).
- After matching, standardized differences were estimated for 367 variables to check for potential residual confusion bias, and those significantly linked to the outcome were use for adjustment in survival analyses
- Cox proportional hazards risk model** were used to compare
 - ✓ The **36-month overall survival** (death)
 - ✓ The **36-month discontinuation free survival** (treatment switch or death)

RESULTS

- In 2014, 3 949 mCRPC patients initiated a 1st-line treatment : 1 162 died during first line and 2 283 had a 2nd-line treatment. Among them :
 - ✓ **693 patients** received the **ABI-DOCE sequence**
 - ✓ **354 patients** received the **DOCE-ABI sequence**
- After trimming and matching: **159 patients per group**




Figure 1. hdPS distribution before and after matching

Table 1. Baseline characteristics at index date before and after matching

	All patients after trimming		Matched patients after trimming		Standardized difference %		
	ABI-DOCE n = 549	DOCE-ABI n = 275	ABI-DOCE n = 159	DOCE-ABI n = 159	Crude	Adjusted	Matched
Median age at index date in years *	73.0	72.0	73.0	73.0	17.3	3.6	10.4
Previous stage of prostate cancer *, %							
mHSPC NDx	15.1	18.5	18.9	18.9	-9.2	1.2	0.0
Progressive mHSPC	14.8	16.4	10.7	10.7	-4.4	0.2	0.0
nmCRPC	17.3	12.7	7.5	7.5	12.8	5.9	0.0
nmHSPC	52.8	52.4	62.9	62.9	0.9	-5.3	0.0
Score de Charlson *					8.0	0.4	3.3
Median [p25% - p75%]	14.0 [13.0;15.0]	14.0 [13.0;15.0]	14.0 [13.0;15.0]	14.0 [13.0;15.0]			
Time since PC diagnosis > 4 years, %	55.0	49.8	45.9	47.8	10.4	-7.1	-3.8
Region of residence of patient, %							
Paris region	14.9	16.0	19.5	15.1	-2.9	1.4	11.7
North-west	22.8	30.9	23.3	32.7	-18.4	-21.8	-21.1
North-east	18.2	19.6	18.2	21.4	-3.6	-5.4	-7.9
South-east	26.8	17.8	26.4	16.4	21.6	23.9	24.7
South-west	15.7	13.1	11.3	12.6	7.3	4.7	-3.9
Overseas territories	1.3	2.5	1.3	1.9	-	-	-

* included in hdPS ; PC = Prostate cancer

Table 2. Description of the three first mCRPC treatment lines

	ABI-DOCE sequence n = 159	DOCE-ABI sequence n = 159
Median duration of 1 st treatment line in months, [p25% - p75%] *	8.4 [4.9;15.4]	6.6 [4.5;9.7]
Median duration of 2 nd treatment line in months, [p25% - p75%] *	6.3 [3.8;8.9]	6.5 [3.1;11.8]
3 rd mCRPC treatment line, %	61.0	59.7
Enzalutamide**	70.1	28.4
Cabazitaxel**	27.8	49.5
Docetaxel**	0.0	17.9
Abiraterone acetate **	1.0	0.0
Combination**	1.0	4.2

*time between first and last infusion for docetaxel and period covered by the dispensed drug for abiraterone **among patients concerned




Figure 2. Sequence of mCRPC treatment lines in matched population in 2014

Table 3. Adjusted overall survival and discontinuation-free survival according to 1st mCRPC treatment line after trimming and matching (Cox model)

	ABI-DOCE sequence n=159	DOCE-ABI sequence n=159	p-value
Overall Survival			
36-month survival probability, % [95%CI]	33.8 [27.4 ; 41.7]	34.4 [27.8 ; 42.5]	0.9105
Median survival time, months [95%CI]	26.2 [22.2 ; 29.8]	26.6 [22.6 ; 30.5]	
Discontinuation-Free Survival			
36-month survival probability, % [95%CI]	13.7 [9.6 ; 19.7]	9.9 [6.4 ; 15.1]	0.1983
Median survival time, months [95%CI]	17.5 [15.4 ; 18.9]	16.1 [14.5 ; 17.6]	




Figure 2. Adjusted overall survival (left) and discontinuation-free survival (right) probability according to the mCRPC treatment sequence after trimming and matching (Cox model)

In real life settings, treatment sequences (ABI-DOCE *versus* DOCE-ABI) seem to have no differential impact on survival outcome in mCRPC patients sharing same characteristics.



Copies of this poster obtained through QR Code are for personal use only and may not be reproduced without permission from ASCO® and the author of this poster.

www.bordeauxpharmacoepi.eu

nicolas.thurin@u-bordeaux.fr

université
de BORDEAUX

CHU
BDX
CENTRE
HOSPITALIER
UNIVERSITAIRE
BORDEAUX

BPE
Bordeaux PharmacoEpi

Inserm

Adera